


Patent claims:

1. The use of a mixture of
A) at least one compound of the formula

5



where R^1 is a C₈-C₂₂ alkyl radical and X^- is an anion, and

- B) a branched fatty acid having a chain length of 8 to 22 carbon atoms
as collector in sylvinite flotation.

10

2. The use as claimed in claim 1, wherein R^1 has a chain length of 12 to 20 carbon atoms.

3. The use as claimed in claim 1 and/or 2, wherein X is a chloride, formate or acetate anion.

15

4. The use as claimed in one or more of claims 1 to 3, wherein constituent A is stearylamine acetate.

20

5. The use as claimed in one or more of claims 1 to 4, wherein constituent B is a branched carboxylic acid having 14 to 22 carbon atoms.

6. The use as claimed in one or more of claims 1 to 5, wherein constituent B is isostearic acid.

25

7. The use as claimed in one or more of claims 1 to 6, wherein the mixing ratio of A:B is 90:10 to 10:90.


8. The use as claimed in one or more of claims 1 to 7, wherein, in addition, glycols, residues from oxoalcohol synthesis and/or water is present.

30

9. The use as claimed in one or more of claims 1 to 8 in amounts of 10 to 500 g/t.

35

10. A composition effective as flotation reagent comprising
A) at least one compound of the formula



5

- where R^1 is a C_8 - C_{22} alkyl radical and X^- is an anion, and
B) a branched fatty acid having a chain length of 8 to 22 carbon atoms,